

An Optical Approach to the Dynamical Casimir Effect

Paweł Węgrzyn

*Marian Smoluchowski Institute of Physics, Jagellonian University, Reymonta 4, 30-059 Cracow, Poland**

We recently proposed a new approach to analyze the parametric resonance in a vibrating cavity based on the analysis of classical optical paths. This approach is used to examine various models of cavities with moving walls. We prove that our method is useful to extract easily basic physical outcome.

PACS numbers: 42.50.Lc, 03.70.+k, 11.10.-z

I. INTRODUCTION

The derivation of the Casimir effect in its original context [1] was the calculation of the electromagnetic force between parallel perfectly conducting plates placed in a vacuum. The geometry of a cavity formed by the parallel plates is also the best context to manifest the nonstationary (dynamical) Casimir effect. The theoretical calculations were first carried out by Moore [2]. The force between plates is dynamically modified, but one encounters more amazing phenomenon being the vacuum emission of photons with nonclassical properties. The latter topic became the subject of intensive studies since it was recognized that the production of photons could be significantly enhanced under the parametric resonance conditions [3, 4]. From the experimental point of view, the hopes for observing vacuum radiation are based on the belief that vibrating cavities can be adjusted to parametric resonance conditions.

Quantum parametric resonance is a complex process and traditional and well-tried methods for calculations in quantum electrodynamics are obsolete here. We must examine open quantum systems under the action of external periodic perturbations. Moreover, the focus of our interest are the instable systems. If we describe physical quantities in resonant systems, then the corresponding mathematical functions inevitably yield narrow peaks or steep stairs. A naive approach to make perturbations in small amplitudes of cavity vibrations breaks down. Therefore, we must go beyond the standard mathematical treatment known from QFT and pursue some new machinery. Most ways of tackling the problem follow the quantization scheme described by Moore [2] in case of one-dimensional cavities. Then, special mathematical techniques are used to trace the evolution of the quantum system with time-dependent boundary conditions. Dodonov and his collaborators [5] reduced the analysis of field equations in a cavity system to some infinite set of coupled and time-dependent oscillators. Another idea based on a concept of renormalization group was proposed in [6, 7]. Our suggestion [8] is to trace the evolution of the quantum fields in a vibrating cavity system following classical optical paths. It is related to the iteration procedure described in [9] and techniques developed for classical field theory in [10] and [11]. In fact, the resonant evolution of the physical system is similar in classical and quantum models. Recently, a similar idea to ground calculations on the analysis of classical optical paths was advocated for a stationary Casimir effect in [12, 13].

In the present paper, we apply our technique [8] to examine cavities with various kinds of their motion and different types of boundary conditions. The calculations are involved in determining the essential physical outcome. In particular, the conditions necessary to keep the parametric resonance and the critical exponents describing resonant evolution are derived. All results are in agreement with those obtained for cavities with moving walls using other techniques. We give more information about resonance windows, more subtle structure of energy densities when amplitudes increase or the adjustment of resonance parameters is getting worse and other new details.

II. A CAVITY WITH ONE OSCILLATING WALL

Most researches on vibrating cavities concentrated on a model of a cavity formed by a static wall and a moving one. The static wall is fixed at $x = 0$, while the moving one follows its trajectory $x = L(t)$. The condition that $L(t) > 0$ is assumed. It is convenient to make use of the billiard function [14]:

$$f(t + L(t)) = t - L(t) . \quad (1)$$

*Electronic address: wegrzyn@th.if.uj.edu.pl

This function contains all information about possible scatterings of massless particles from the moving wall. It can be also understood as the trajectory of the wall in light-cone coordinates. Its derivatives are recognized as retarded Doppler factors (the retardation relation is $t = t^* + L(t^*)$):

$$\dot{f}(t) = \frac{1 - \dot{L}(t^*)}{1 + \dot{L}(t^*)} . \quad (2)$$

Consider now a classical scalar field $A(t, x)$ in a one-dimensional cavity. It obeys the wave equation:

$$(\partial_t^2 - \partial_x^2)A(t, x) = 0 . \quad (3)$$

The field is subject to some boundary conditions. Usually, we assume either Dirichlet's boundary conditions:

$$A(t, 0) = A(t, L(t)) = 0, \quad (4)$$

or Neumann's boundary conditions at the instantaneous positions of walls:

$$\partial_x A(t, 0) = (\dot{L}(t)\partial_t - \partial_x)A(t, L(t)) = 0, \quad (5)$$

In both cases, the classical solutions are given by:

$$A(t, x) = \varphi(t + x) \mp \varphi(t - x), \quad (6)$$

where the upper sign corresponds to Dirichlet's case, while the lower one to Neumann's case. The profile function is subject to the following relation:

$$\varphi(\tau) = \varphi(f(\tau)) . \quad (7)$$

The energy density of the classical wave packet and its total energy are given by:

$$T_{00}(t, x) = 1/2(\partial_t A)^2 + 1/2(\partial_x A)^2 = \varrho(t + x) + \varrho(t - x) , \quad (8)$$

$$E(t) = \int_0^{L(t)} dx T_{00}(t, x) = \int_{t-L(t)}^{t+L(t)} d\tau \varrho(\tau) , \quad (9)$$

where $\varrho(\tau) = \dot{\varphi}^2(\tau)$. Let us now describe classical optical paths. We consider a massless particle that starts at time τ moving right from the position $x = 0$. Denote by $T_n(\tau)$ and $T_n^*(\tau)$ times of n -th collisions of the particle with the moving wall and the static one respectively. It is easy to notice that $T_n(\tau) = (f^{-1})^{\circ n}(\tau)$, where the notation $(f^{-1})^{\circ n}(\tau)$ means here and throughout this paper n -fold composition $f^{-1} \circ f^{-1} \circ \dots \circ f^{-1}$, while $T_n^*(\tau)$ are to be determined from the retardation relations $T_n^*(\tau) + L(T_n^*(\tau)) = T_n(\tau)$. The parametric resonance is related to the existence of periodic particle trajectories [8, 11]. Each periodic trajectory obeys a following condition (for any non-negative integer n):

$$T_n(\tau_0) = \tau_0 + nT , \quad (10)$$

where τ_0 is a starting point in time and T is a period. The retarded times are respectively $T_n^*(\tau_0) = T_n(\tau_0) - T/2$. Any periodic particle trajectory appears on condition that the mirror trajectory has return points:

$$L(\tau_0^* + nT) = T/2 . \quad (11)$$

It is usually assumed that $T/2$ refers to the length of the static cavity, so that $T/2 \equiv L \equiv L(0)$. The crucial characteristic of a particle trajectory is the cumulative Doppler factor [8]:

$$D_n(\tau) = \frac{1}{T_n(\tau)} = \prod_{k=1}^n \dot{f}(T_k(\tau)) = \prod_{k=1}^n \frac{1 - \dot{L}(T_k^*(\tau))}{1 + \dot{L}(T_k^*(\tau))} . \quad (12)$$

The above function has a direct physical interpretation. If a massless particle starts at time τ moving right inside the cavity, then after n rebounds from both cavity walls its kinetic energy changes by the factor $D_n(\tau)$. A trajectory is called positive (or negative respectively) if its cumulative Doppler factor tends to infinity (or zero) with increasing n . We denote by τ_+ and τ_- starting points for positive and negative periodic trajectories. If positive trajectories

exist, then any trajectory approaches one of the positive trajectories for long times. The same is true for negative trajectories if we consider the evolution backward in time. In other words, $T_n(\tau) \rightarrow T_n(\tau_+)$ and $T_n^{-1}(\tau) \rightarrow T_n^{-1}(\tau_-)$ for large n .

The evolution of the energy density can be deduced from the following formula,

$$\varrho(T_n(\tau)) = \varrho(\tau) D_n^2(\tau) , \quad (13)$$

together with the asymptotic approximation for long times (i.e. $n \gg 1$) [8]:

$$\varrho(nT + \tau_+ + \varepsilon D_n^{-1}(\tau_+)) \cong \varrho(\tau_+ + \varepsilon) D_n^2(\tau_+ + \varepsilon) , \quad (14)$$

$$\varrho(nT + \tau_- + \varepsilon) \cong \varrho(\tau_- + \varepsilon D_n(\tau_-)) D_n^2(\tau_- + \varepsilon D_n(\tau_-)) . \quad (15)$$

The above formulas explain the formation of travelling narrow packets in the energy density $T_{00}(t, x)$ [3, 5, 15]. The profile function $\varrho(\tau)$ has peaks located at the spots of positive periodic trajectories τ_+ . The height of a peak grows like $D_n^2(\tau_+)$, while its width diminishes like $D_n^{-1}(\tau_+)$. The value of the energy density near the peak decreases like $D_n^2(\tau_-)$. The evolution of the total energy can be analyzed using the formula (we assume here $\tau_0 \leq L$):

$$E(T_n^*(\tau_0)) = \int_{T_{n-1}(\tau_0)}^{T_n(\tau_0)} d\tau \varrho(\tau) = \int_{f(\tau_0)}^{\tau_0} d\tau \varrho(\tau) D_n(\tau) . \quad (16)$$

To calculate the total accumulated energy inside the cavity after some time we need only an initial energy density (i.e. $\varrho(\tau)$ for $\tau \leq L$) and the knowledge of cumulative Doppler factors. In fact, it is usually enough to know these factors in the vicinities of spots of periodic trajectories.

Now, we turn into the basic model of vibrating cavities specified by the sinusoidal cavity wall motion:

$$L(t) = L + \Delta L \sin(\omega t) , \quad (17)$$

where $\Delta L < L$ and $\omega \Delta L < 1$. The parametric resonance frequencies are $\omega_N = N\pi/L$, where N is the order of the resonance. Positive and negative periodic particle trajectories correspond to the following starting points:

$$\tau_{+m} = \frac{(-N + 2m + 1)L}{N} , \quad \tau_{-m} = \frac{(-N + 2m)L}{N} , \quad (18)$$

where $m = 0, 1, \dots, N - 1$. The cumulative Doppler factors can be calculated:

$$\begin{aligned} D_n(\tau_{\pm m}) &= \left(\frac{1 \pm \omega_N \Delta L}{1 \mp \omega_N \Delta L} \right)^n \\ &\cong \exp(\pm 2n\omega_N \Delta L) \quad \text{for } \Delta L \ll L . \end{aligned} \quad (19)$$

We conclude that there are N travelling peaks in the energy density. The average distances between peaks are equal. For N even, $N/2$ peaks are moving right and $N/2$ peaks are moving left. For odd N the number of peaks of each kind differs by one. All the peaks have the same height and the same width. The total energy grows exponentially with time. These classical results agree with the results of the corresponding quantized cavity model [4]. This fact will be explained later.

The above picture is true provided that the amplitude of cavity oscillations is small. Strictly speaking, we need $\Delta L/L < N$. For larger amplitudes, the behavior of the cavity system is slightly modified. If $\Delta L/L > M/N$ ($M = 1, 2, 3, \dots$), then there appear additional positive trajectories corresponding to the following return points (11):

$$\omega_N \Delta L \sin(\omega_N \tau_{M\pm}^*) = M\pi , \quad (20)$$

and the corresponding cumulative Doppler factors are:

$$D_n(\tau_{M\pm}) = \left(\frac{1 \pm \sqrt{(\omega_N \Delta L)^2 - (M\pi)^2}}{1 \mp \sqrt{(\omega_N \Delta L)^2 - (M\pi)^2}} \right)^n . \quad (21)$$

It follows that beside the first series of N peaks in the energy density, there appear next M series of N peaks each. The peaks of additional series are smaller and located between higher peaks.

For practical purposes, it is important to consider off resonant behavior of the cavity system. The instability in the field theory appears not only for finely tuned frequencies. We are usually dealing with some band structure. To establish the resonance window, we move the resonant frequency: $\omega = \omega_N + \Delta\omega$. The equation for return points (11) is now:

$$\sin(\omega\tau^*) = -\frac{L\Delta\omega}{\omega\Delta L}, \quad (22)$$

so that solutions exist provided that:

$$\frac{\Delta\omega}{\omega} < \frac{\Delta L}{L}. \quad (23)$$

This condition agrees with the result of the quantized model [16]. It defines the windows for the resonant frequencies to keep the resonant instability. The cumulative Doppler factors at starting points of periodic trajectories are:

$$D_n(\tau_{\pm m}) = \left(\frac{1 \pm \sqrt{(\omega\Delta L)^2 - (L\Delta\omega)^2}}{1 \mp \sqrt{(\omega\Delta L)^2 - (L\Delta\omega)^2}} \right)^n. \quad (24)$$

Again, the above picture is exact provided that $\Delta L/L + (1 + 1/N)\Delta\omega/\omega > 1/N$. Otherwise, next series of smaller peaks in the energy density will appear.

Now, we present quantum formulas. The vacuum expectation value of the energy density is given by [2]:

$$\langle T_{00}(t, x) \rangle = \varrho(t+x) + \varrho(t-x), \quad (25)$$

where

$$\varrho(\tau) = -\frac{\pi}{48}\dot{R}^2(\tau) - \frac{1}{24\pi}S[R](\tau), \quad (26)$$

and the Schwartz derivative is defined by:

$$S[R] = \frac{\ddot{R}}{\dot{R}} - \frac{3}{2} \left(\frac{\ddot{R}}{\dot{R}} \right)^2. \quad (27)$$

The function $R(\tau)$ is subject to Moore's equation:

$$R(\tau) - R(f(\tau)) = 2, \quad (28)$$

where our billiard function appears in the quantum case as well. The formula for the evolution of the energy density is given by:

$$\varrho(T_n(\tau)) = \varrho(\tau)D_n^2(\tau) + A_n(\tau)D_n^2(\tau), \quad (29)$$

where the cumulative Doppler factor is defined again by the same formula Eq. (12) and the definition of the quantum term called the cumulative conformal anomaly contribution [8] is given below:

$$A_n(\tau) = \frac{1}{24\pi}S[T_n](\tau) = -\frac{1}{24\pi} \sum_{k=1}^n D_k^{-2}(\tau)S[f](T_k(\tau)). \quad (30)$$

The quantum formula for the total energy accumulated in the cavity is:

$$\begin{aligned} E(T_n^*(\tau_0)) &= \int_{T_{n-1}(\tau_0)}^{T_n(\tau_0)} d\tau \varrho(\tau) \\ &= \int_{f(\tau_0)}^{\tau_0} d\tau [\varrho(\tau) + A_n(\tau)] D_n(\tau). \end{aligned} \quad (31)$$

The quantum anomaly implies the difference between classical and quantum formulas for the evolution of the energy. However, under the resonance conditions the leading role is played by the cumulative Doppler factor. This fact explains why most of classical results match quantum ones. It refers to the band structure of resonance frequencies, the formation and the shape of travelling packets in the energy density, the exponential growth of the total energy. To make predictions for these features, the classical theory could be good enough. But it is not true for the lowest resonance channel. In this case, the anomalous mechanism of energy growth clashes with the resonant enhancement of the initial vacuum fluctuations [8]. The lack of the instability for the lowest resonance frequency is also related to the $SL(2, R)$ symmetry of the cavity system [17, 18], which is the remainder of the conformal group.

Finally, let us also conclude from the above results that there is no difference between imposing Dirichlet's and Neumann's boundary conditions in the context of our cavity models.

III. A CAVITY WITH TWO OSCILLATING WALLS

A natural generalization of the considerations of the previous section is to allow both walls to oscillate [5, 7, 19]. Consider now that the right cavity wall moves along the trajectory $x = L_1(t)$, while the left one follows $x = -L_2(t)$. We assume also that the cavity was static in the past, so that $L_1(t) = L_2(t) = L/2$ for $t < 0$, and L corresponds to the static cavity length. The general formulas given in this section will be rigorously justified if we assure that $L_1(t) > 0$ and $L_2(t) > 0$ (the walls never collides) and the wall velocities are never close to the velocity of light. The billiard functions are implemented for both trajectories ($i = 1, 2$):

$$f_i(t + L_i(t)) = t - L_i(t) . \quad (32)$$

Moreover, it is convenient to define an additional pair of billiard functions:

$$f_L = f_2 \circ f_1 \quad , \quad f_R = f_1 \circ f_2 . \quad (33)$$

They describe double reflections of massless particles from the cavity walls. The set of classical optical paths is now more complex. Consider again a massless particle that starts at time τ moving from the position $x = 0$. Denote by $T_{Ln}(\tau)$ (and $T_{Rn}(\tau)$ respectively) times when the particle returns to its initial position $x = 0$ after n rebounds from both walls assuming that the first collision is with the left (or with the right) wall. It is easy to find that $T_{L(R)n}(\tau) = (f_{L(R)}^{-1})^{\circ n}(\tau)$. Further, $T_{L(R)n}^*(\tau)$ and $T_{L(R)n}^{**}(\tau)$ are times of successive collisions with the left (right) and the right (left) walls for both types of trajectories respectively. They can be derived from the following relations:

$$T_{L(R)n}^*(\tau) + L_{1(2)}(T_{L(R)n}^*(\tau)) = T_{L(R)n}(\tau) , \quad (34)$$

$$T_{L(R)n}^{**}(\tau) + L_{2(1)}(T_{L(R)n}^{**}(\tau)) = f_{1(2)}(T_{L(R)n}(\tau)) . \quad (35)$$

For both types of trajectories cumulative Doppler factors are defined:

$$\begin{aligned} D_{L(R)n}(\tau) &= \frac{1}{\tilde{T}_{L(R)n}(\tau)} = \prod_{k=1}^n \dot{f}_{L(R)}(T_{L(R)k}(\tau)) \\ &= \prod_{k=1}^n \frac{1 - \dot{L}_{1(2)}(T_{L(R)k}^*(\tau))}{1 + \dot{L}_{1(2)}(T_{L(R)k}^*(\tau))} \prod_{k=1}^n \frac{1 - \dot{L}_{2(1)}(T_{L(R)k}^{**}(\tau))}{1 + \dot{L}_{2(1)}(T_{L(R)k}^{**}(\tau))} . \end{aligned} \quad (36)$$

Classical solutions of a wave equation inside a cavity with two moving walls are given by:

$$A(t, x) = \varphi_L(t + x) \mp \varphi_R(t - x), \quad (37)$$

where again the upper sign corresponds to Dirichlet's boundary conditions and the lower one to Neumann's boundary conditions on the walls. The profile functions in both cases obey the same relations:

$$\varphi_L(\tau) = \varphi_R(f_1(\tau)) , \quad \varphi_R(\tau) = \varphi_L(f_2(\tau)) . \quad (38)$$

The energy density of the classical wave packet and its total energy can be represented as:

$$T_{00}(t, x) = \varrho_L(t + x) + \varrho_R(t - x) , \quad \varrho_{L(R)}(\tau) = \dot{\varphi}_{L(R)}^2(\tau) , \quad (39)$$

$$E(t) = \int_{-L_2(t)}^{L_1(t)} dx T_{00}(t, x) = \int_{t-L_2(t)}^{t+L_1(t)} d\tau \varrho_L(\tau) + \int_{t-L_1(t)}^{t+L_2(t)} d\tau \varrho_R(\tau) . \quad (40)$$

The classical evolution of the system can be traced with the help of the following formulas:

$$\varrho_{L(R)}(T_{L(R)n}(\tau)) = \varrho_{L(R)}(\tau) D_{L(R)n}^2(\tau) , \quad (41)$$

$$E(t) = \int_{f_2(\tau_2)}^{\tau_1} d\tau \varrho_L(\tau) D_{Ln}(\tau) + \int_{f_1(\tau_1)}^{\tau_2} d\tau \varrho_R(\tau) D_{Rn}(\tau) , \quad (42)$$

where $\tau_{1(2)} \equiv f_{L(R)}^{\circ n}(t + L_{1(2)}(t))$. If we take n big enough to support that $\tau_{1(2)} \leq L/2$, then we obtain the formula that allows us to calculate the evolution of the total energy from the knowledge of Doppler factors and the energy distribution in the initial state.

Before we turn into examples of cavity systems, we explain why many specific systems of cavities with two oscillating walls just reproduce results similar to those obtained in cavity models with one oscillating wall. The reason is that billiard functions (33) correspond to some physical trajectories in two dimensions. It means that we can find a trajectory $x = L(t)$ which describes some realistic motion of the wall such that:

$$f_{L(R)}(t + L(t)) = t - L(t) . \quad (43)$$

We can verify that $L(t) > 0$, $L(t) = L$ for $t < 0$ and $\dot{L}(t)$ is never close to the speed of light. Obviously, in general such "effective" trajectories are different for f_L and f_R , but in both cases the following composition rules are satisfied:

$$L(t) = L_1(t_1) + L_2(t_2) , \quad (44)$$

$$\frac{1 - \dot{L}(t)}{1 + \dot{L}(t)} = \frac{1 - \dot{L}_1(t_1)}{1 + \dot{L}_1(t_1)} \frac{1 - \dot{L}_2(t_2)}{1 + \dot{L}_2(t_2)} , \quad (45)$$

where $t \pm L(t) = t_1 \pm L_1(t_1)$ and $t_1 \mp L_1(t_1) = t_2 \pm L_2(t_2)$ for f_L (the upper sign) and f_R respectively.

The latter relation Eq.(45) states just that $\dot{L}(t)$ is the relativistic sum of velocities $\dot{L}_1(t_1)$ and $\dot{L}_2(t_2)$. However, this composition rule has nothing to do with any relative motion since there exists no inertial frame where we could make respective time moments synchronous. Nevertheless, both left- and right-movers in cavities with two oscillating walls correspond to some models of cavities with a single oscillating wall. In other words, each mode of a cavity system with two moving walls can be described by a simpler model discussed in the previous section.

Let us discuss specific types of symmetric and antisymmetric oscillations of cavities. In the paper of Dalvit and Mazzitelli [7], these types of cavity oscillations are called breathing modes ("electromagnetic antishaker") and translational modes ("electromagnetic shaker"). For symmetric oscillations we have $L_1(t) = L_2(t)$ and the cavity oscillates symmetrically with respect to the center of the cavity. The billiard functions and Doppler factors are identical, namely $f_1 \equiv f_2$, $f_L \equiv f_R$ and $D_L \equiv D_R$. The model is equivalent to the cavity model with one oscillating wall, where the prescribed trajectory of the wall is defined by the billiard function $f_1 \circ f_1$. For antisymmetric oscillations we have $L_1(t) + L_2(t) = L$, and the cavity is oscillating as a whole with its length kept constant in the laboratory frame. Then, $f_{1(2)}(\tau) = f_{2(1)}^{-1}(\tau - L) - L$ and $f_{L(R)}(\tau) = f_{1(2)}^{-1}(f_{1(2)}(\tau) - L) - L$. It is interesting to note that for periodic translational cavity oscillations with L being a period, the billiard functions f_L and f_R correspond to static ones. It follows that there is no effect of the cavity movement on the energy distribution of fields and the whole cavity system is equivalent to the static one.

Finally, we analyze harmonic oscillations of cavity walls:

$$\begin{aligned} L_1(t) &= \frac{L}{2} + \Delta L_1 \sin(\omega_R t) , \\ L_2(t) &= \frac{L}{2} + \Delta L_2 \sin(\omega_L t - \delta) + \Delta L_2 \sin \delta . \end{aligned} \quad (46)$$

Again, we want to establish resonance windows, find the evolution of the total energy and describe the shape of the local energy density. We must first detect positive and negative periodic trajectories of massless particles moving inside the cavity. They can be derived from the generalization of Eq.(11):

$$L_1(t_1) + L_2(t_1 \pm \frac{T}{2}) = \frac{T}{2} , \quad (47)$$

where the upper (lower) sign corresponds to the "left" T_{L_n} ("right" T_{R_n}) trajectories. The cumulative Doppler factors at the starting points of periodic trajectories can be calculated then:

$$\begin{aligned} D_{L(R)n} &= \left(\frac{1 - \omega_R \Delta L_1 \cos(\omega_R t_1)}{1 + \omega_R \Delta L_1 \cos(\omega_R t_1)} \right)^n \left(\frac{1 - \omega_L \Delta L_2 \cos(\omega_L t_1 \mp \omega_L T/2 - \delta)}{1 + \omega_L \Delta L_2 \cos(\omega_L t_1 \mp \omega_L T/2 - \delta)} \right)^n \\ &\cong \exp(-2n[\omega_R \Delta L_1 \cos(\omega_R t_1) + \omega_L \Delta L_2 \cos(\omega_L t_1 \mp \omega_L T/2 - \delta)]) \quad \text{for } \Delta L_i \ll L . \end{aligned} \quad (48)$$

First, consider resonant oscillations with $\omega_L = \omega_R = \omega_N \equiv N\pi/L$. If there is no dephasing $\delta = 0$, the system is equivalent to the cavity with one oscillating wall. The oscillation frequency is ω_N , the effective cavity length is L and

the effective amplitude of oscillations is $\Delta L_1 + (-1)^N \Delta L_2$. For $\Delta L_1 = \Delta L_2$ and N odd the effective cavity system is static. The same is true for $\Delta L_1 = \Delta L_2$, $\delta = \pi$ and N even. These results agree with the paper [7]. If the amplitudes of oscillations are equal $\Delta L_1 = \Delta L_2 \equiv \Delta L$ and $\delta \neq 0, \pi$, then the solutions of Eq.(47) are $t_1 = (\delta L/\pi + 2k)L/N$ and $t_1 = (N - 2k - 1)L/N$, where $k = 0, 1, \dots, N - 1$. The respective cumulative Doppler factors are:

$$D_n(\tau_{k\pm}) \cong \exp(\pm 2n\omega_N \Delta L(1 + (-1)^N \cos \delta)) . \quad (49)$$

There are N positive and N negative particle trajectories. It follows that there are N peaks in the energy density. The calculations for $\Delta L_1 \neq \Delta L_2$ are a bit more complicated. However, if $\Delta L_1 + (-1)^N \Delta L_2 \cos \delta \neq 0$ then there exist N positive and N negative particle trajectories. The corresponding cumulative Doppler factors are given by:

$$D_n \cong \exp(\pm 2n\omega_N (\Delta L_1 + (-1)^N \Delta L_2 \cos \delta)) . \quad (50)$$

For $\Delta L_1 + (-1)^N \Delta L_2 \cos \delta = 0$ there is no exponential instability in the cavity system.

To establish the resonance window we assume $\omega_L = \omega_R \equiv \omega_N + \Delta\omega$. The relation Eq.(47) yields:

$$\Delta L_1 \sin(\omega t_1) + (-1)^N \Delta L_2 \sin(\omega t_1 - \delta) + \Delta L_2 \sin \delta = \frac{-L\Delta\omega}{\omega} . \quad (51)$$

To keep the parametric resonance conditions for the vibrating cavity Eq.(46) it is enough to support that:

$$\frac{\Delta\omega}{\omega} < \frac{\Delta L_2 \sin \delta + \sqrt{(\Delta L_1)^2 + (\Delta L_2)^2 + 2(-1)^N \Delta L_1 \Delta L_2 \cos \delta}}{L} . \quad (52)$$

The quantum formulas for cavities with two oscillating walls include anomaly contributions in the respective way. Since the modifications are straightforward, we skip their discussion here.

IV. CONCLUSIONS

An optical approach to the dynamical Casimir effect based on the construction of classical optical paths is useful. In particular, it is a powerful technique to analyze resonance systems. We can trace the behavior of a cavity system with moving walls when we increase amplitudes or loose fine tuning for resonance eigenfrequencies. We can calculate modified critical exponents, find more subtle structure of travelling peaks in energy densities and control the conditions for the instability of the system. We have established some "composition laws" coming into effect if we want a more complicated model of a cavity with two moving walls reduce to the analysis of a cavity with one moving wall. Moreover, our understanding of the mechanism of parametric resonance in cavities with moving walls and the implications of quantum field theory is much better. For example, we can judge when some semi-classical approximation is accurate or we can explain why some resonances are not observed at the quantum level. It is further remarkable that we can distinguish between exponential and power-like instability and observe the role of symmetries at the quantum level [18].

We have seen in several examples that many qualitative or even quantitative conclusions can be drawn from classical versions of respective cavity models. Presumably, the parametric resonance mechanism and evolution laws are similar in both cases. However, we should be careful about quantum anomalies that can significantly modify our predictions. A subtle role of quantum modifications can be traced within our approach.

The results derived for Dirichlet's and Neumann's boundary conditions for one-dimensional cavities are the same. This point was already made in [20]. The cavities with two oscillating walls possess a double set of characteristic functions, namely two families of both cumulative Doppler factors and cumulative anomaly contributions. It is related to the fact that from a given space-time point there are two independent directions to follow trajectories of massless particles. In three dimensions, such functions depend also on localizations and orientations.

-
- [1] H.B.G. Casimir, Proc. Kon. Ned. Akad. Wetensch. **51**, 793 (1948).
 - [2] G.T. Moore, J. Math. Phys. **11**, 2679 (1970).
 - [3] C.K.Law, Phys. Rev. Lett. **73**, 1931 (1994).
 - [4] V.V.Dodonov, Phys. Lett. A **207**, 126 (1995).
 - [5] V.V.Dodonov, Modern Nonlinear Optics, Part 1, p.309-394
 - [6] D.A. Dalvit, F.D. Mazzitelli, Phys. Rev. A **57**, 2113 (1998).

- [7] D. A. Dalvit, F.D. Mazzitelli, Phys. Rev. A **59**, 3049 (1999) .
- [8] P. Węgrzyn, Phys. Lett. A **322**, 263 (2004).
- [9] C.K. Cole, W.C. Schieve, Phys.Rev. **A 52**, 4405 (1995).
- [10] J.Dittrich, P.Duclos, P.Šeba, Phys. Rev.**E 49**, 3535 (1994).
- [11] O. Méplan, C. Gignoux, Phys. Rev. Lett. **76**, 408 (1996).
- [12] R.L. Jaffe, A. Scardicchio, Phys. Rev. Lett. **92**, 070402 (2004).
- [13] R.L. Jaffe, A. Scardicchio, arXiv: quant-ph/0406041 (2004).
- [14] P. Węgrzyn, T. Róg, Acta Phys. Polon. **32**, 129 (2001).
- [15] A.Lambrecht, M.T.Jaeckel, S.Reynaud, Phys. Rev. Lett. **77**, 615 (1996).
- [16] P. Węgrzyn, T. Róg, Acta Phys. Polon. **B 34**, 3887 (2003).
- [17] Y.Wu, X.Yang, J. Phys. **A 34**, 327 (2001).
- [18] P.Węgrzyn, Mod. Phys. Lett. **A 19**, 769 (2004).
- [19] J.Y.Ji, H.H.Jung, K.S.Soh, Phys.Rev. **A57**, 4952 (1998).
- [20] D.T.Alves, C.Farina, P.A.Maia Neto, J. Phys. **A 36**, 11333 (2003).